

This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike License](https://creativecommons.org/licenses/by-nc-sa/4.0/). Your use of this material constitutes acceptance of that license and the conditions of use of materials on this site.



Copyright 2009, The Johns Hopkins University and John McGready. All rights reserved. Use of these materials permitted only in accordance with license rights granted. Materials provided "AS IS"; no representations or warranties provided. User assumes all responsibility for use, and all liability related thereto, and must independently review all materials for accuracy and efficacy. May contain materials owned by others. User is responsible for obtaining permissions for use from third parties as needed.



JOHNS HOPKINS
BLOOMBERG
SCHOOL *of* PUBLIC HEALTH

Comparing Proportions between Two Independent Populations

John McGready
Johns Hopkins University

Lecture Topics

- Using CIs for difference in proportions between two independent populations
- Large sample methods for comparing proportions between two populations
 - Normal method
 - Chi-squared test
- Fisher's exact test
- Relative risk



JOHNS HOPKINS
BLOOMBERG
SCHOOL *of* PUBLIC HEALTH

Section A

The Two Sample z-Test for Comparing Proportions
between Two Independent Populations: The Confidence
Interval Approach

Comparing Two Proportions

- We will motivate by using data from the Pediatric AIDS Clinical Trial Group (ACTG) Protocol 076 Study Group*
- Study design
 - “We conducted a randomized, double-blinded, placebo-controlled trial of the efficacy and safety of zidovudine (AZT) in reducing the risk of maternal-infant HIV transmission”
 - 363 HIV infected pregnant women were randomized to AZT or placebo

Notes: *Conner, E., et al. (1994). Reduction of maternal-infant transmission of human immunodeficiency virus type 1 with zidovudine treatment, *New England Journal of Medicine* 331: 18.

Comparing Two Proportions

- Results
 - Of the 180 women randomized to AZT group, 13 gave birth to children who tested positive for HIV within 18 months of birth
 - Of the 183 women randomized to the placebo group, 40 gave birth to children who tested positive for HIV within 18 months of birth

Notes on Design

- Random assignment of Tx
 - Helps insure two groups are comparable
 - Patient and physician could not request particular treatment
- Double blind
 - Patient and physician did not know treatment assignment

Observed HIV Transmission Proportions

- AZT

$$\hat{p}_{AZT} = \frac{13}{180} = 0.07 = 7\%$$

- Placebo

$$\hat{p}_{PLAC} = \frac{40}{183} = 0.22 = 22\%$$

HIV Transmission Proportions: 95% CIs

```
. cii 180 13
```

Variable	Obs	Mean	Std. Err.	-- Binomial Exact -- [95% Conf. Interval]	
	180	.0722222	.019294	.0390137	.1203358

```
. cii 183 40
```

Variable	Obs	Mean	Std. Err.	-- Binomial Exact -- [95% Conf. Interval]	
	183	.2185792	.0305507	.160984	.2855248

Notes on HIV Transmission Proportions

- Is the difference significant, or can it be explained by chance?
- Since CIs do not overlap suggests significant difference
 - Can we compute a confidence interval on the difference in proportions?
 - Can we compute a p-value?

Sampling Distribution: Difference in Sample Proportions


- Since we have large samples we know the sampling distributions of the sample proportions in both groups are approximately normal
- It turns out the difference of quantities, which are (approximately) normally distributed, are also normally distributed

Sampling Distribution: Difference in Sample Proportions

- So, the big news is . . .
 - The sampling distribution of the difference of two sample proportions, each based on large samples, approximates a normal distribution
 - This sampling distribution is centered at the true (population) difference, $p_1 - p_2$


Sampling Distribution: AZT Group

- Simulated sampling distribution of sample proportion: AZT group




Sampling Distribution: Difference in Sample Proportions

- Simulated sampling distribution of sample proportion: placebo group



Sampling Distribution: Difference in Sample Proportions

- Simulated sampling distribution of difference in sample proportions:
AZT - placebo



95% Confidence Interval for Difference in Proportions

- Our most general formula

$$\text{best estimate from sample} \pm 2 \times SE(\text{best estimate from sample})$$

- The best estimate of a population difference based on sample proportions:

$$\hat{p}_1 - \hat{p}_2$$

- Here, \hat{p}_1 may represent the sample proportion of infants HIV positive (within 18 months of birth) for 180 infants in the AZT group, and \hat{p}_2 may represent the sample proportion of infants HIV positive (within 18 months of birth) for 183 infants in the AZT group

95% CI for Difference in Proportions: AZT Study

- So, $\hat{p}_1 - \hat{p}_2 = 0.07 - 0.22 = -0.15$: hence the formula for the 95% CI for $p_1 - p_2$ is:

$$-0.15 \pm 2 \times SE(\hat{p}_1 - \hat{p}_2)$$

- Where $SE(\hat{p}_1 - \hat{p}_2)$ = standard error of the difference of two sample proportions

Standard Error of Difference in Proportions

- Statisticians have developed formulas for the standard error of the difference
- These formulas depend on sample sizes in both groups and sample proportions in both groups
- The $SE(\hat{p}_1 - \hat{p}_2)$ is greater than either $SE(\hat{p}_1)$ or $SE(\hat{p}_2)$
 - Why do you think this is?

Principle

- Variation from independent sources can be added
 - Why do you think this is additive?

$$SE(\hat{p}_1 - \hat{p}_2) = \sqrt{\frac{p_1 \times (1 - p_1)}{n_1} + \frac{p_2 \times (1 - p_2)}{n_2}}$$

- Of course, we don't know p_1 and p_2 : so we estimate with \hat{p}_1 and \hat{p}_2 to get an estimated standard error:

$$\hat{SE}(\hat{p}_1 - \hat{p}_2) = \sqrt{\frac{\hat{p}_1 \times (1 - \hat{p}_1)}{n_1} + \frac{\hat{p}_2 \times (1 - \hat{p}_2)}{n_2}}$$

Comparing Two Independent Groups: HIV/AZT Study

- Recall the data from the Infant HIV/ AZT study

	Group	
	AZT	Placebo
Number of subjects (n)	64	68
Proportion Infants HIV+ Within 18 Months	0.07	0.22

$$SE(\hat{p}_1 - \hat{p}_2) = \sqrt{\frac{.07 \times .93}{180} + \frac{.22 \times .78}{183}} \approx .36$$

95% CI for Difference in Proportions: HIV/ AZT Study

- So in this example, the estimated 95% for the true difference in proportions of infants contracting HIV between the AZT and placebo groups:

$$-0.15 \pm 2 \times SE(\hat{p}_1 - \hat{p}_2)$$

$$-0.15 \pm 2 \times .036$$

$$-0.15 \pm .072$$

$$-0.222 \text{ to } 0.078 \approx$$

$$-22\% \text{ to } -8\%$$

Summary: AZT Study

■ Results

- The proportion of infants who tested positive for HIV within 18 months of birth was seven percent (95% CI 4 -12%) in the AZT group and twenty-two percent in the placebo group (95% CI 16 - 28%)
- The study results estimate the absolute decrease in the proportion of HIV positive infants born to HIV positive mothers associated with AZT to be as low as 8% and as high as 22%