


This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike License](https://creativecommons.org/licenses/by-nc-sa/4.0/). Your use of this material constitutes acceptance of that license and the conditions of use of materials on this site.



Copyright 2008, The Johns Hopkins University and Brian Caffo. All rights reserved. Use of these materials permitted only in accordance with license rights granted. Materials provided "AS IS"; no representations or warranties provided. User assumes all responsibility for use, and all liability related thereto, and must independently review all materials for accuracy and efficacy. May contain materials owned by others. User is responsible for obtaining permissions for use from third parties as needed.

Lecture 20

Brian Caffo

Department of Biostatistics
Johns Hopkins Bloomberg School of Public Health
Johns Hopkins University

November 6, 2007

Table of contents

- 1 Table of contents
- 2 Outline
- 3 Recap
- 4 The delta method
- 5 Derivation of the delta method

- 1 Review two sample binomial results
- 2 Delta method

Two sample binomials results

Recall $X \sim \text{Bin}(n_1, p_1)$ and $Y \sim \text{Bin}(n_2, p_2)$. Also this information is often arranged in a 2×2 table:

$n_{11} = x$	$n_{12} = n_1 - x$	n_1
$n_{21} = y$	$n_{22} = n_2 - y$	n_2

- $\hat{RD} = \hat{p}_1 - \hat{p}_2$

$$\hat{SE}_{\hat{RD}} = \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_2(1-\hat{p}_2)}{n_2}}$$

- $\hat{RR} = \frac{\hat{p}_1}{\hat{p}_2}$

$$\hat{SE}_{\log \hat{RR}} = \sqrt{\frac{(1-\hat{p}_1)}{\hat{p}_1 n_1} + \frac{(1-\hat{p}_2)}{\hat{p}_2 n_2}}$$

- $\hat{OR} = \frac{\hat{p}_1/(1-\hat{p}_1)}{\hat{p}_2/(1-\hat{p}_2)} = \frac{n_{11}n_{22}}{n_{12}n_{21}}$

$$\hat{SE}_{\log \hat{OR}} = \sqrt{\frac{1}{n_{11}} + \frac{1}{n_{12}} + \frac{1}{n_{21}} + \frac{1}{n_{22}}}$$

CI = Estimate $\pm Z_{1-\alpha/2} \text{SE}_{\text{Est}}$

Standard errors

Table of contents

Outline

Recap

The delta method

Derivation of the delta method

- **delta method** can be used to obtain large sample standard errors
- Formally, the delta method states that if

$$\frac{\hat{\theta} - \theta}{\hat{SE}_{\hat{\theta}}} \rightarrow N(0, 1)$$

then

$$\frac{f(\hat{\theta}) - f(\theta)}{f'(\hat{\theta})\hat{SE}_{\hat{\theta}}} \rightarrow N(0, 1)$$

- Asymptotic mean of $f(\hat{\theta})$ is $f(\theta)$
- Asymptotic standard error of $f(\hat{\theta})$ can be estimated with $f'(\hat{\theta})\hat{SE}_{\hat{\theta}}$

Example

- $\theta = p_1$
- $\hat{\theta} = \hat{p}_1$
- $\hat{SE}_{\hat{\theta}} = \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1}}$
- $f(x) = \log(x)$
- $f'(x) = 1/x$
- $\frac{\hat{\theta} - \theta}{\hat{SE}_{\hat{\theta}}} \rightarrow N(0, 1)$ by the CLT
- Then $\hat{SE}_{\log \hat{p}_1} = f'(\hat{\theta})\hat{SE}_{\hat{\theta}}$

$$= \frac{1}{\hat{p}_1} \sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n_1}} = \sqrt{\frac{(1-\hat{p}_1)}{\hat{p}_1 n_1}}$$

- And

$$\frac{\log \hat{p}_1 - \log p_1}{\sqrt{\frac{(1-\hat{p}_1)}{\hat{p}_1 n_1}}} \rightarrow N(0, 1)$$

Putting it all together

- Asymptotic standard error

$$\begin{aligned}\text{Var}(\log \hat{R}R) &= \text{Var}\{\log(\hat{p}_1/\hat{p}_2)\} \\ &= \text{Var}(\log \hat{p}_1) + \text{Var}(\log \hat{p}_2) \\ &\approx \frac{(1 - \hat{p}_1)}{\hat{p}_1 n_1} + \frac{(1 - \hat{p}_2)}{\hat{p}_2 n_2}\end{aligned}$$

- The last line following from the delta method
- The approximation requires large sample sizes
- The delta method can be used similarly for the log odds ratio

Motivation for the delta method

Table of
contents

Outline

Recap

The delta
methodDerivation of
the delta
method

- If $\hat{\theta}$ is close to θ then

$$\frac{f(\hat{\theta}) - f(\theta)}{\hat{\theta} - \theta} \approx f'(\hat{\theta})$$

- So

$$\frac{f(\hat{\theta}) - f(\theta)}{f'(\hat{\theta})} \approx \hat{\theta} - \theta$$

- Therefore

$$\frac{f(\hat{\theta}) - f(\theta)}{f'(\hat{\theta})\hat{SE}_{\hat{\theta}}} \approx \frac{\hat{\theta} - \theta}{\hat{SE}_{\hat{\theta}}}$$