


This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike License](https://creativecommons.org/licenses/by-nc-sa/4.0/). Your use of this material constitutes acceptance of that license and the conditions of use of materials on this site.



Copyright 2009, The Johns Hopkins University and John McGready. All rights reserved. Use of these materials permitted only in accordance with license rights granted. Materials provided "AS IS"; no representations or warranties provided. User assumes all responsibility for use, and all liability related thereto, and must independently review all materials for accuracy and efficacy. May contain materials owned by others. User is responsible for obtaining permissions for use from third parties as needed.



JOHNS HOPKINS
BLOOMBERG
SCHOOL *of* PUBLIC HEALTH

Lecture 2b: Practice Problems

John McGready
Johns Hopkins University

Practice Problems

- Suppose a population is normally distributed (and you are a member of the population)
 1. If you have a standard score of $Z = 2$, what percentage of the population has scores greater than you?
 2. If you have a standard score of $Z = -2$, what percentage of the population has scores greater than you?
 3. If you have a standard score of $Z = 1$, what percentage of the population has scores less than you?
 4. If you have a standard score of $Z = 1$, what percentage of the population has scores farther away from the population mean (in either direction) than you?
 5. If you have a standard score of $Z = -1.7$, what percentage of the population has scores farther away from the population mean (in either direction) than you?
 6. If you have a standard score of $Z = -1.7$, what percentage of the population has scores greater than you?

Practice Problems

7. Suppose the distribution of grades in your statistics class is normal, with mean = 83.4, $s = 7$. There are 120 total in the class. If you score a 97.4 in the class, roughly how many people have scores higher than you?
8. Suppose we call unusual observations in a population of normally distributed values those that are either at least 2 SD above the mean or about 2 SD below the mean. What percent are unusual? In other words, what percent of the observations will have a standard score either $Z > + 2.0$ or $Z < -2.0$? What percent would have $|Z| > 2$?