


This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike License](https://creativecommons.org/licenses/by-nc-sa/4.0/). Your use of this material constitutes acceptance of that license and the conditions of use of materials on this site.



Copyright 2009, The Johns Hopkins University and John McGready. All rights reserved. Use of these materials permitted only in accordance with license rights granted. Materials provided "AS IS"; no representations or warranties provided. User assumes all responsibility for use, and all liability related thereto, and must independently review all materials for accuracy and efficacy. May contain materials owned by others. User is responsible for obtaining permissions for use from third parties as needed.



JOHNS HOPKINS
BLOOMBERG
SCHOOL *of* PUBLIC HEALTH

Lecture 6a: Practice Problems

John McGready
Johns Hopkins University

Practice Problems

- A study was performed on a representative sample of 258 intravenous drug users (IVDUs)
- Of particular interest to the researchers were factors which may influence the risk of contracting tuberculosis amongst IVDUs*
- Ninety-seven of the study subjects admitted to sharing needles to shoot drugs
- Of these ninety-seven, twenty-four had a positive tuberculin test result
- The other 161 subjects denied having shared needles—of these 161 subjects, 28 had a positive tuberculin test result

Notes: * Based on data reported in Graham, N., et al. Prevalence of tuberculin positivity and skin test anergy in HIV-1-seropositive and seronegative intravenous drug users, *Journal of the American Medical Association* 267: 3.

Practice Problems

- Using these study results, construct a 95% confidence interval for the difference in the proportion of tuberculosis infected IVDUS who shared needles as compared to IVDUS who did not share needles