


This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike License](https://creativecommons.org/licenses/by-nc-sa/4.0/). Your use of this material constitutes acceptance of that license and the conditions of use of materials on this site.



Copyright 2009, The Johns Hopkins University and John McGready. All rights reserved. Use of these materials permitted only in accordance with license rights granted. Materials provided "AS IS"; no representations or warranties provided. User assumes all responsibility for use, and all liability related thereto, and must independently review all materials for accuracy and efficacy. May contain materials owned by others. User is responsible for obtaining permissions for use from third parties as needed.



JOHNS HOPKINS  
BLOOMBERG  
SCHOOL *of* PUBLIC HEALTH

## Section B

---

Two Sample t-test: Getting a p-value

# Hypothesis Test to Compare Two Independent Groups

- Two sample (unpaired) t-test
- Is the (mean) weight change equal in the two diet groups?
  - $H_0: \mu_1 = \mu_2$
  - $H_A: \mu_1 \neq \mu_2$
- In other words, is the expected difference in weight change zero?
  - $H_0: \mu_1 - \mu_2 = 0$
  - $H_A: \mu_1 - \mu_2 \neq 0$

# Hypothesis Test to Compare Two Independent Groups

- Recall, general “recipe” for hypothesis testing . . .
  1. Start by assuming  $H_0$  true
  2. Measure distance of sample result from  $\mu_0$  (here again its 0)
  3. Compare test statistic (distance) to appropriate distribution to get p-value

$$t = \frac{(\text{observed dif f}) - (\text{null dif f})}{SE \text{ of observed dif f erence}}$$

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\hat{SE}(\bar{x}_1 - \bar{x}_2)} = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

## Diet Type and Weight Loss Study

- In the diet types and weight loss study, recall:

$$\bar{x}_1 - \bar{x}_2 = -3.9kg$$

$$SE(\bar{x}_1 - \bar{x}_2) = 1.17kg$$

- So in this study:

$$t = \frac{-3.9}{1.17} \approx -3.3$$


- So this study result was 3.3 standard errors below the null mean of 0 (i.e., 3.3 standard errors from the mean weight less expected if null was true)

## How Are p-values Calculated?

- Is a result 3.3 standard errors below 0 unusual?
  - It depends on what kind of distribution we are dealing with
- The p-value is the probability of getting a test statistic as extreme as (or more extreme than) what you observed (-3.3) by chance if was true
- The p-value comes from the sampling distribution of the difference in two sample means
- What is the sampling distribution of the difference in sample means?
  - If both groups are large (more than 60 subjects) then this distribution is approximately normal
  - This sampling distribution will be centered at true difference
  - Under null hypothesis, this true difference is 0

# Diet/Weight Loss Sample

- To compute a p-value, we would need to compute the probability of being 3.3 or more standard errors away from 0 on a standard normal curve



# How to Use Stata to Perform a 2-Sample T-Test

- Command syntax:

- `ttesti  $n_1$   $\bar{x}_1$   $s_1$   $n_2$   $\bar{x}_2$   $s_2$ , unequal`

```
. ttesti 64 -5.7 8.6 68 -1.8 3.9, unequal
```

```
Two-sample t test with unequal variances
```

	Obs	Mean	Std. Err.	Std. Dev.	[ 95% Conf. Interval]	
x	64	-5.7	1.075	8.6	-7.848216	-3.551784
y	68	-1.8	.4729445	3.9	-2.744001	-.8559989
combined	132	-3.690909	.5978226	6.868458	-4.873545	-2.508273
diff		-3.9	1.174437		-6.234436	-1.565564

```
diff = mean(x) - mean(y)                                t = -3.3207  
Ho: diff = 0                                           Satterthwaite's degrees of freedom = 86.6941
```

```
Ha: diff < 0  
Pr(T < t) = 0.0007
```

```
Ha: diff != 0  
Pr(|T| > |t|) = 0.0013
```

```
Ha: diff > 0  
Pr(T > t) = 0.9993
```

# How to Use Stata to Perform a 2-Sample T-Test

- Command syntax:
  - `ttesti  $n_1$   $\bar{x}_1$   $s_1$   $n_2$   $\bar{x}_2$   $s_2$ , unequal`

```
. ttesti 64 -5.7 8.6 68 -1.8 3.9, unequal
```

```
Two-sample t test with unequal variances
```

	Obs	Mean	Std. Err.	Std. Dev.	[ 95% Conf. Interval]	
x	64	-5.7	1.075	8.6	-7.848216	-3.551784
y	68	-1.8	.4729445	3.9	-2.744001	-.8559989
combined	132	-3.690909	.5978226	6.868458	-4.873545	-2.508273
diff		-3.9	1.174437		-6.234436	-1.565564

```
diff = mean(x) - mean(y)                                t = -3.3207
Ho: diff = 0                                             Satterthwaite's degrees of freedom = 86.6941

Ha: diff < 0                                           Ha: diff != 0                                           Ha: diff > 0
Pr(T < t) = 0.0007                                     Pr(|T| > |t|) = 0.0013                                   Pr(T > t) = 0.9993
```

# How to Use Stata to Perform a 2-Sample T-Test

- Command syntax:
  - `ttesti  $n_1$   $\bar{x}_1$   $s_1$   $n_2$   $\bar{x}_2$   $s_2$ , unequal`

```
. ttesti 64 -5.7 8.6 68 -1.8 3.9, unequal
```

```
Two-sample t test with unequal variances
```

	Obs	Mean	Std. Err.	Std. Dev.	[ 95% Conf. Interval]
x	64	-5.7	1.075	8.6	-7.848216 -3.551784
y	68	-1.8	.4729445	3.9	-2.744001 -.8559989
combined	132	-3.690909	.5978226	6.868458	-4.873545 -2.508273
diff		-3.9	1.174437		-6.234436 -1.565564

```
diff = mean(x) - mean(y) t = -3.3207  
Ho: diff = 0 Satterthwaite's degrees of freedom = 86.6941
```

```
Ha: diff < 0  
Pr(T < t) = 0.0007
```

```
Ha: diff != 0  
Pr(|T| > |t|) = 0.0013
```

```
Ha: diff > 0  
Pr(T > t) = 0.9993
```

# Summary: Weight Loss Example

## ■ Statistical method

- “We randomly assigned 132 severely obese patients . . . to a carbohydrate restricted (low-carbohydrate) diet or a calorie- and fat-restricted diet”
- “For comparison of continuous variables between the two groups, we calculated the change from baseline to six months in each subject, and compared the mean changes in the two diet groups using an unpaired t-test”

## ■ Result

- “Subjects on the low-carbohydrate diet lost more weight than those on a low fat diet (95% confidence interval for the difference in weight loss between groups, -1.6 to -6.2 kg;  $p < .01$ )”