


This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike License](https://creativecommons.org/licenses/by-nc-sa/4.0/). Your use of this material constitutes acceptance of that license and the conditions of use of materials on this site.



Copyright 2009, The Johns Hopkins University and John McGready. All rights reserved. Use of these materials permitted only in accordance with license rights granted. Materials provided "AS IS"; no representations or warranties provided. User assumes all responsibility for use, and all liability related thereto, and must independently review all materials for accuracy and efficacy. May contain materials owned by others. User is responsible for obtaining permissions for use from third parties as needed.




JOHNS HOPKINS
BLOOMBERG
SCHOOL *of* PUBLIC HEALTH

Lecture 2a: Practice Problem Solutions

John McGready
Johns Hopkins University

Practice Problems

2. Consider a large, randomized study such as the Salk polio vaccine trial



Practice Problems


2. How does randomization help to ensure that the outcome /treatment group relationship observed is not confounded by any factors known or unknown? More specifically, what part of the necessary conditions for confounding does randomization eliminate?

Practice Problems

- Remember, a necessary condition for a variable to be a confounder is that it must be related to both the outcome and the predictor of interest
- In randomized studies, the “treatment” is the predictor of interest
- Randomization with a large number of potential subjects eliminates the association between the predictor of interest and any other variable besides (potentially) the outcome
- In other words, randomization allows the only difference between the “treatment” groups to be the treatment of interest


Practice Problems

- In the “confounding conditions” diagram with outcome **Y**, predictor of interest **X**, and potential confounder **Z**:



Practice Problems

- Randomization eliminates the necessary link between X and Z



Practice Problems

3. Suppose a study were performed to assess the relationship between a diet choice (vegan, lacto-ovo vegetarian, neither) and cholesterol level. Subjects were not randomized to a dietary group. Before interpreting the relationship, can you name some potential factors for which it would be advisable to control?

Solutions

- Remember, for a factor to be a confounder, it must be associated with both cholesterol level and with being vegetarian

Solutions

- Some candidates include the following:
 - Sex—females are more likely to be vegetarian and have lower cholesterol
 - Smoking—vegetarians are less likely to smoke and smoking is associated with higher cholesterol
 - Income—vegetarians are wealthier and increased wealth associated with lower cholesterol