

This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike License](https://creativecommons.org/licenses/by-nc-sa/4.0/). Your use of this material constitutes acceptance of that license and the conditions of use of materials on this site.



Copyright 2006, The Johns Hopkins University and Karl Broman. All rights reserved. Use of these materials permitted only in accordance with license rights granted. Materials provided "AS IS"; no representations or warranties provided. User assumes all responsibility for use, and all liability related thereto, and must independently review all materials for accuracy and efficacy. May contain materials owned by others. User is responsible for obtaining permissions for use from third parties as needed.

## Statistics for laboratory scientists

### Solutions for the homework problems for lecture 18

1. We use the following R code to do this:

```
x <- c(55.2, 58.1, 41.7, 44.9, 44.8, 48.9, 47.5,  
      48.1, 48.4, 51.6, 40.6, 48.0)  
y <- c(48.7, 52.6, 65.2, 70.4, 44.2, 54.7, 44.0,  
      66.5, 56.8)  
wilcox.test(x,y)
```

This gives a P-value of 8.2% (compare this with the t-test, which gave  $P = 5.1\%$ ).

---

[ [3rd term syllabus](#) | [4rd term syllabus](#) | [R for Windows](#) ]

Last modified: Thu Mar 2 12:57:31 EST  
2006