

This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike License](https://creativecommons.org/licenses/by-nc-sa/4.0/). Your use of this material constitutes acceptance of that license and the conditions of use of materials on this site.



Copyright 2009, The Johns Hopkins University and John McGready. All rights reserved. Use of these materials permitted only in accordance with license rights granted. Materials provided "AS IS"; no representations or warranties provided. User assumes all responsibility for use, and all liability related thereto, and must independently review all materials for accuracy and efficacy. May contain materials owned by others. User is responsible for obtaining permissions for use from third parties as needed.



JOHNS HOPKINS  
BLOOMBERG  
SCHOOL *of* PUBLIC HEALTH

## Lecture 6c: Practice Problems

---

John McGready  
Johns Hopkins University

## Example: Wages and Education Level

1. Recall equation of regression line relating estimated mean hourly wages (U.S. \$, 1985) to years of education: from Stata . . .

$$\hat{y} = -0.75 + 0.75x$$

- a) What is the estimated mean hourly wage (in 1985) for persons with 12 years of education?
- b) What is the estimated difference in hourly wages (in 1985) for persons with 16 years of education versus 12 years of education?

## Example: Arm Circumference and Sex

2. Recall the regression relating arm circumference to child's sex for the random sample of 150 Nepali children less than 12 months old

$$\hat{y} = 12.5 + -0.13x$$

- In this example,  $x$  is the binary variable for sex, coded as a 1 for female children and 0 for male children; suppose  $x$  was coded as 1 for male children and 0 for female children
  - a) What would the resulting slope estimate be?
  - b) What would the resulting intercept estimate be?