


This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike License](https://creativecommons.org/licenses/by-nc-sa/4.0/). Your use of this material constitutes acceptance of that license and the conditions of use of materials on this site.



Copyright 2008, The Johns Hopkins University and Brian Caffo. All rights reserved. Use of these materials permitted only in accordance with license rights granted. Materials provided "AS IS"; no representations or warranties provided. User assumes all responsibility for use, and all liability related thereto, and must independently review all materials for accuracy and efficacy. May contain materials owned by others. User is responsible for obtaining permissions for use from third parties as needed.

Lecture 27

Brian Caffo

Department of Biostatistics
Johns Hopkins Bloomberg School of Public Health
Johns Hopkins University

December 13, 2007

Table of contents

① Table of contents

② Outline

③ Multiplicity

④ Bonferoni

⑤ FDR

Outline

- 1 Familywise error rates
- 2 Bonferoni procedure
- 3 Performance of Bonferoni with multiple independent tests
- 4 False discovery rate procedure

Multiplicity

- After rejecting a χ^2 omnibus test you do all pairwise comparisons
- You conducted a study with 20 outcomes and 30 different combinations of covariates. You consider significance at all combinations.
- You compare diseased tissue versus normal tissue expression levels for 20k genes
- You compare rest versus active at 300k voxels in an fMRI study

Multiplicity

- Performing two α -level tests:

H_0^1 versus H_a^1 and H_0^2 versus H_a^2

E_1 Reject H_0^1 and E_2 Reject H_0^2

$$\begin{aligned}
 FWE &= P(\text{one or more false rej} \mid H_0^1, H_0^2) \\
 &= P(E_1 \cup E_2 \mid H_0^1, H_0^2) \\
 &= P(E_1 \mid H_0^1, H_0^2) + P(E_2 \mid H_0^1, H_0^2) \\
 &\quad - P(E_1 \cap E_2 \mid H_0^1, H_0^2) \\
 &\leq P(E_1 \mid H_0^1, H_0^2) + P(E_2 \mid H_0^1, H_0^2) \\
 &= 2 \times \alpha
 \end{aligned}$$

Result : The **familywise error rate** for k hypotheses tested at level α is bounded by $k\alpha$

E_i - false rejection for test i

All probabilities are conditional on all of the nulls being true

$$\begin{aligned} FWE &= P(\text{one or more false rej}) \\ &= P(\cup_{i=1}^k E_i) \\ &= P\left\{E_1 \cup (\cup_{i=2}^k E_i)\right\} \\ &\leq P(E_1) + P(\cup_{i=2}^k E_i) \\ &\quad \vdots \\ &\leq P(E_1) + P(E_2) + \dots + P(E_k) \\ &= k\alpha \end{aligned}$$

Other direction

- The *FWE* is no larger than $k\alpha$ where k is the number of tests
- The *FWE* is no smaller than α

$$P(\cup_{i=1}^k E_i) \geq P(E_1) = \alpha$$

- The lower bound is obtained when the E_i are identical
 $E_1 = E_2 = \dots = E_k$
- **Bonferroni's** tests each individual hypothesis at level $\alpha^* = \alpha/k$
 - The *FWE* is no larger than $k\alpha^* = k\alpha/k = \alpha$
 - The *FWE* is no smaller than α/k

Bonferroni's procedure

If α^* is small and the tests are independent, then the upper bound on the *FWE* is nearly obtained

$$\begin{aligned} FWE &= P(\text{one or more false rej}) \\ &= 1 - P(\text{no false rej}) \\ &= 1 - P(\cap_{i=1}^k \bar{E}_i) \\ &= 1 - (1 - \alpha^*)^k \\ &\approx 1 - (1 - k\alpha^*) \\ &= k\alpha^* = \alpha \end{aligned}$$

Recall the approximation for α^* near 0

$$\frac{f(\alpha^*) - f(0)}{\alpha^* - 0} \approx f'(0)$$

hence

$$f(\alpha^*) \approx f(0) + \alpha^* f'(0)$$

In our case $f(\alpha^*) = (1 - \alpha^*)^k$ so $f(0) = 1$

$f'(\alpha^*) = -k(1 - \alpha^*)^{k-1}$ so $f'(0) = -k$

Therefore $(1 - \alpha^*)^k \approx 1 - k\alpha^*$

- For Bonferoni's procedure $\alpha^* = \alpha/k$ so will be close to 0 for a large number of tests
- When there are lots of tests that are (close to) independent, the upper bound on the *FWE* used is appropriate
- When the test are closely related, then the *FWE* will be closer to the lower bound, and Bonferoni's procedure is conservative
- Is the familywise error rate always the most appropriate quantity to control for?

- The **false discovery rate** is the proportion of tests that are falsely declared significant
- Controlling the FDR is less conservative than controlling the FWE rate
- Introduced by Benjamini and Hochberg

Benjamini and Hochberg procedure

- 1 Order your k p-values, say $p_1 < p_2 < \dots < p_k$
- 2 Define $q_i = kp_i/i$
- 3 Define $F_i = \min(q_i, \dots, q_k)$
- 4 Reject for all i so that F_i is less than the desired FDR

Note that the F_i are increasing, so you only need to find the largest one so that $F_i < FDR$

1st 10 of 50 SNPs (Rosner page 581)

Gene	i	p_i	$q_i = kp_i/i$	F_i
30	1	<.0001	.0035	.0035
20	2	.011	.28	.16
48	3	.017	.28	.16
50	4	.017	.22	.16
4	5	.018	.18	.16
40	6	.019	.16	.16
7	7	.026	.18	.18
14	8	.034	.21	.21
26	9	.042	.23	.23
47	10	.048	.24	.24

Example

- Bonferoni cutoff $.05/50 = .001$; only the first Gene is significant
- For a FDR of 0 – 15%; only the first Gene would be declared significant
- For a FDR of 16 – 20%, the first 7 would be significant